

Biron Plus

Iron, Zinc & Folic acid

COMPOSITION

Biron Plus Capsule: Each capsule contains Dried Ferrous Sulphate BP 150 mg, Zinc Sulphate Monohydrate USP 61.80 mg and Folic Acid BP 0.50 mg.

DESCRIPTION

Biron Plus is the combined preparation of carbonyl iron, folic acid and zinc. It contains carbonyl iron with not less than 98% iron content. Carbonyl iron, having high bioavailability and low toxicity is safer and more effective choice for iron supplementation.

INDICATIONS

It is indicated for the treatment and prophylaxis of iron, folic acid and zinc deficiency especially during pregnancy and lactation.

DOSAGE AND ADMINISTRATION

Adult: one capsule daily before food or as directed by the physician.

CONTRAINDICATION

It is contraindicated in patients with known hypersensitivity to any of its component or those with iron overload.

USE IN PREGNANCY AND LACTATION

Use of any drug during the first trimester of pregnancy should be avoided if possible. Thus administration of iron during the first trimester requires definite evidence of iron deficiency. Prophylaxis of iron deficiency where inadequate diet calls for supplementary zinc and folic acid is justified during the remainder of pregnancy.

PRECAUTIONS

Special care should be taken in patient with iron overload states, such as haemochromatosis, haemolytic anaemia or red cell aplasia. Failure to response to the treatment requires further investigations to exclude other causes of anaemia. In patients with renal failure there may be the risk of zinc accumulation.

SIDE EFFECTS

Gastrointestinal irritations such as nausea, anorexia, vomiting, discomfort, constipation and diarrhoea may occur. Patients may complain of dark stool. Carbonyl iron pellets are incorporated into the capsules to reduce the possibility of gastrointestinal irritations. Rarely there may be allergic reactions.

OVERDOSE

Symptoms of carbonyl iron include decreased energy, nausea, abdominal pain, tarry stool; a weak, rapid pulse, fever; coma; seizures.

DRUG INTERACTIONS

Carbonyl iron may decrease the absorption of tetracycline antibiotics, quinolone antibiotics, levodopa, levothyroxine, methyl dopa and penicillamine. Folic acid interacts with antiepileptics, so plasma concentrations of phenobarbital, phenytoin and primidone are possibly reduced.

STORAGE

Store in a cool & dry place (Below 25° C). Keep away from light. Keep out of reach of children.

COMMERCIAL PACK

Biron Plus Capsule: Each box containing 6X10 TR capsules in blister pack.

Manufactured by



For further query on the use of this medicine, consult to a registered Doctor or Pharmacist.