

Myoclop

Clopidogrel USP 75mg

COMPOSITION

Myoclop 75 mg Tablet : Each film coated tablet contains Clopidogrel Bisulfate USP equivalent to Clopidogrel 75 mg .

PHARMACOLOGY

Myoclop (Clopidogrel) is a selective ADP receptor antagonist to inhibit platelet aggregation. Myoclop (Clopidogrel) selectively inhibits the binding of Adenosine diphosphate (ADP) to its platelet receptor and the subsequent ADP-mediated activation of the glycoprotein IIb/IIIa complex, thereby inhibiting platelet aggregation.

INDICATION

Myoclop is indicated for the prevention of atherosclerotic events in patients with history of symptomatic atherosclerotic disease (ischemic stroke, myocardial infarction or established peripheral arterial disease). It is given prophylactically as an alternative to aspirin in patients at risk of thrombo-embolic disorders such as myocardial infarction and stroke.

DOSAGE AND ADMINISTRATION

The recommended dose of Myoclop 75 mg is once daily, with or without food. No dosage adjustment is necessary for elderly patients or patients with renal disease.

CONTRAINDICATION AND PRECAUTION

Clopidogrel is contraindicated in the following conditions:

- Hypersensitivity to the drug substance or any component of the product.
- Active pathological bleeding such as peptic ulcer or intracranial hemorrhage.

SIDE EFFECT

Clopidogrel is generally well tolerated drug. A few common side-effects like hemorrhage (including gastro-intestinal and intracranial), abdominal discomfort, nausea, vomiting, diarrhoea, headache, dizziness, vertigo, paraesthesia, rash, pruritus, hepatic and biliary disorder, neutropenia may occur.

DRUG INTERACTION

Drugs that might induce lesions with a propensity to bleed (e.g. ulcer) like aspirin, NSAIDs, should be used with cautions to patients taking Clopidogrel. No significant interactions were observed in the co-administration with atenolol, nifedipine or both of the drugs. Cautions should be used in co-administration of phenytoin, tamoxifen, tolbutamide, warfarin, fluvastatin with Clopidogrel for possible impaired metabolism.

USE IN PREGNANCY AND LACTATION

There are no adequate & well-controlled studies of Clopidogrel in pregnant women. So Clopidogrel should be used in pregnancy only if clearly needed. Clopidogrel should not be prescribed to nursing mother unless absolutely necessary.

STORAGE CONDITION

Store below 30°C. Keep all medicines out of the reach of children.

HOW SUPPLIED

Myoclop 75 mg Tablet : Box containing 3 x 10 tablets in blister pack.

Manufactured by



For further query on the use of this medicine, consult to a registered Doctor or Pharmacist.