

Tolpa

Tolperisone Hydrochloride

Presentation

Tolpa 50 mg Tablet: Each film coated tablet contains Tolperisone Hydrochloride INN 50 mg.

Description

Tolperisone Hydrochloride is a centrally acting muscle relaxant. It effectively reduces Muscular hypertonicity, improves peripheral blood flow by vasodilatation and reduces pain.

Indications and uses

Increased tone of skeletal muscles due to organic neurological disorders (e.g. injury of the pyramidal tract, multiple sclerosis, myelopathy, encephalomyelitis etc.), muscular hypertension, muscular spasm, muscular contracture, rigidity, spinal automatism and discopathy. Obliterative vascular diseases (obliterative arteriosclerosis, diabetic angiopathy, obliterative thromboangitis, Raynaud's disease, diffuse scleroderma) and disorders due to injured innervation of the vessels (acrocyanosis, intermittent angioneurotic dysbasia). In individual cases post-thrombotic venous and lymphatic circulation disorders, crural ulcer.

Dosage and Administration

One should follow doctor's instructions regarding the dosage of Tolpa. The Recommended dose is as followings:

Adult: 50-150 mg three times daily according to the individual requirement.

Contraindications

Hypersensitivity to any component of the product

Myasthenia gravis

Nursing mother

Precautions

In case of children, the prescribed dose and duration of treatment should closely be observed.

Side-effects

Sometimes, muscular weakness, headache, nausea, vomiting and rarely hypersensitivity reactions may occur as side effects.

Use in Pregnancy and Lactation

Tolperisone hydrochloride should be used in pregnancy according to physician's advice. Tolperisone hydrochloride should not be used during breast feeding.

Drug Interactions

Concomitant use of methocarbamol with tolperisone hydrochloride has been reported to Cause disturbance of visual accommodation.

Over dosage

In the event of over dosage supportive therapy may be substituted.

Commercial Pack

Tolpa 50 mg Tablet: Each box contains 5 blister strips of 10 tablets.

Manufactured by



For further query on the use of this medicine, consult to a registered Doctor or Pharmacist.